


**Clinical Updates**

## Keppra IVPB changing to IV Push: Go-live 1/8/26


- Keppra IV Push administration guidelines will be added to the comments
- Plans Impacted
  - Severe Traumatic Brain Injury (GCS < 9) Subplan (MSJ)
  - NS Neurosurgery Cranial Surgery Plan (MSJ)
  - Neurology Subarachnoid Hemorrhage (SAH) Plan (MSJ)
  - Mild to Moderate Traumatic Brain Injury Subplan, Adult (GCS 9-15)
  - Seizure Treatment Plan, Adult (System)
  - ED Seizures Plan (System)
  - APA Subarachnoid Hemorrhage (MSJ)
  - FSER Seizures Plan

Seizure Treatment	Current State
	 <b>Loading Dose</b>  PROVIDER: consider discontinuing Keppra before 7 days if aneurysm is secured and the patient has not had seizure events.
<input type="checkbox"/>    levETIRacetam (Keppra)	20 mg/kg, IVPB, IV Piggyback, ONCE For seizure treatment
<input type="checkbox"/> +12 hr   levETIRacetam (Keppra)	1000, mg, IVPB, IV Piggyback, BID, Duration: 7 day
<input checked="" type="checkbox"/> Non Categorized	

Seizure Treatment	Future State
	 <b>Loading Dose</b>  PROVIDER: consider discontinuing Keppra before 7 days if aneurysm is secured and the patient has not had seizure events.
<input type="checkbox"/>    levETIRacetam (Keppra)	20 mg/kg, Inj, IV Push, ONCE For seizure treatment. Push over 3-5 min. Max dose of 4500 mg IVP over 5 min. Pharmacist to round to the nearest 500 mg. May be administered undiluted.
<input type="checkbox"/> +12 hr   levETIRacetam (Keppra)	1000, mg, Inj, IV Push, BID, Duration: 7 day Push over 3-5 min. May be administered undiluted.

## New Smart Zone Alerts for Positive Flu and Measles

To promote patient and caregiver safety, a new Smart Zone alert will fire when a patient has a positive Flu or a Positive Measles result.



Page 2 of 2