
Clinical Reminders

GO LIVE
NOV 5

Independent Double Check Documentation

- Additional medications will be added to the High Alert Medication Independent Double Check (IDC) documentation requirements.
- ***In addition to* the current IDC requirements, medications that will prompt a co-sign in the MAR will now include:**
  - IV insulin
  - IV opioid infusions
  - Neuromuscular blocking agents
  - Hypertonic solutions (>0.9% NaCl)
  - IV push sedative agents (Ketamine, Phenobarbital)
  - Argatroban
  - TPN
  - Population Specific Requirements Remain
    - Mannitol (Pediatrics only)
    - Concentrated electrolytes (Potassium Chloride) – Pediatrics and NICU only
    - All controlled substances (NICU only)
    - Antiarrhythmics (NICU only)

Follow required IDC process and obtain a second RN co-sign in the EMR:



Refer to 1MM.MED.0001 "High Alert Medications and Independent Double Check (IDC) – Adults and Pediatrics" policy

Clinical Reminders

GO LIVE
NOV 5

Additional Adult and Pediatric Code Cart Changes

ADULT
- Dopamine Premix - Remove
- D50W Syringe - Add 1 additional syringe

PEDIATRIC
- 4.2% Infant Bicarb - remove from Peds Code Cart Tray
- 8.4% bicarb - Add 1, 50 mEq syringe
- Calcium gluconate - remove from Peds Code Cart Tray
- Calcium chloride - Add 2, 1 gm syringes
- Pediatric Code Sheet - Add
- Dopamine Premix - remove from Peds Code Cart Tray
- D50W Syringe - Add 1 additional syringe

Clinical Reminders

ROLLING DIVISION
GO-LIVE

BD Alaris™ Infusion System Updates

PEDIATRICS • Full system: Point of Care Unit (PCU) / Pump / Syringe / Patient Controlled Analgesia (PCA) / $EtCO_2$

RED DOTS MUST MATCH — Point of Care Unit (PCU) + EVERY ATTACHED MODULE
Use updated red-dot components together only. Never attach an unmarked older module to an updated PCU.

CORRECT • All components marked




INCORRECT • Mixed marked/unmarked



New Programming Screens


DELAY FOR

"Delay Until" is removed. Use "Delay For" (1 min to 11 hr 59 min); verify the programmed duration.



CUSTOM CONCENTRATION: MORE INFO

A two-step Guardrails alert shows the entered drug amount, diluent volume and concentration. Verify before proceeding.



- HIGH-PRIORITY ALARMS: "Infusion Complete" and "Infusion Complete–KVO" now require immediate response.
- $EtCO_2$: "NO CO_2" replaces "No Breath." Assess patient and sampling setup; message meaning is not an apnea diagnosis.
- SYRINGE / PCA SUPPLIES: Check manufacturer, model, size, flow rate and current approved disposables before loading.
- CLEAN + DISINFECT: Clean first, then disinfect with a BD-approved product and required contact/drying time. Never clean while attached to a patient. Include PCA handset/cord.

30-SECOND SAFETY CHECK: Red dots match • Correct patient/profile • Guardrails/settings verified • Syringe/sets approved

Clinical Reminders

ROLLING DIVISION
GO-LIVE

BD Alaris™ Infusion System Updates

ADULT ACUTE CARE • PCA USERS ONLY

RED DOTS MUST MATCH — Point of Care Unit (PCU) + EVERY ATTACHED MODULE
Use updated red-dot components together only. Never attach an unmarked older module to an updated PCU.

CORRECT • All components marked




INCORRECT • Mixed marked/unmarked



PCA / $EtCO_2$ Screen Examples


PCA CUSTOM CONCENTRATION

Two-step Guardrails warning: select MORE INFO and verify drug amount, diluent volume and concentration.



$EtCO_2$: "NO CO_2" (IF USED)

"No Breath" becomes "No CO_2." Assess the patient and sampling setup; this text does not diagnose apnea.



PCA Syringes • Examples from BD v12.6.X compatible list

SIZE	BD-COMPATIBLE PCA SYRINGE / MODEL
20 mL	BD 302830
30 mL	BD 302832; IMS Pump-Jet® NDC 76329-1912-1
35 mL	Cardinal Health Monoject™ detachable-plunger barrel 8881135609
50 mL	BD 309653

Clinical Reminders

EDUCATION BOOST

Insulin Answers! Rapid Facts

Meal & Correction Insulin

Goal: Insulin given within 30 minutes of CBG and meal tray arrival.






Safety pearls: Do not give MEAL insulin if the tray is delayed or the patient is not eating. Give CORRECTION insulin regardless of NPO. Reassess CBG if the meal is delayed >30 minutes. “HOVER” over MAR orders for rapid-acting insulin dose accuracy.

Clinical Reminders

EDUCATION BOOST
WOUND & SKIN

PIP Tip: Wound Photos

iMobile photo workflow



- **Photos taken within the app DO NOT upload to the EMR**
- **Complex wounds, new pressure injuries, PI stage 3 and above: notify wound team about photos**

Photos taken within the app DO NOT upload to the EMR. For complex wounds, new pressure injuries, and pressure injuries stage 3 and above, notify the wound team about photos.

Clinical Reminders


GO LIVE NOV 9

MOD Retirement - Accessing the New Medication Calculator Links


Cerner build update

- MOD is being retired.
- The "Clinical Resources" button in the toolbar is being removed and replaced with links to (1) LexiDrug and (2) the new SharePoint Med Calculator page.
- Users can continue to access Med Dosing Calculators via PowerPlan links and Reference Text links; those links are being updated to point to the new SharePoint med calculator link.


CURRENT STATE - Toolbar




FUTURE STATE - Toolbar



FUTURE STATE - SharePoint Med Calculators



NOT CHANGING - PowerPlan Access



Not changing: PowerPlans and Reference Text continue to be accessed the same way; only the link endpoint is changing.